

Herding behavior of agents as a background of financial fluctuations

Aleksejus Kononovicius, V. Gontis, B. Kaulakys

Institute of Theoretical Physics and Astronomy, Vilnius University
aleksejus.kononovicius@gmail.com

2012-07-09

Financial fluctuations





Figure: Statistical properties of the absolute one minute returns in the financial markets: (a) probability and (b) spectral density. Similar statistical properties are also observed for the trading activity.

Frequent large deviations (fat tails), long-range memory (power law spectral density and thus, due to W-K theorem, auto-correlation).

The wild side of the financial markets




Experiment by Deneubourg



Taken from [Detrain & Deneubourg, 2006 (Physics of Life Reviews)].

Kirman's formulation of herding model



X dynamics

are determined by the one-step transition probabilities:

$$\begin{aligned} p(X \rightarrow X + 1) &= (N - X)\sigma_1 + hX(N - X), \\ p(X \rightarrow X - 1) &= X\sigma_2 + hX(N - X). \end{aligned}$$

Statistical properties of Kirman's herding model





Figure: Probability density functions of $x = X/N$ in the Kirman's herding model: (a) $\sigma = \sigma_1 = \sigma_2$, (b) $\sigma_1 \neq \sigma_2$. (a) $\sigma < h$ (red curve), $\sigma = h$ (blue curve), $\sigma > h$ (magenta curve). (b) $h > \sigma_1 > \sigma_2$ (red curve), $h < \sigma_1 < \sigma_2$ (blue curve).

Model of human behavior? (restaurant scenario)



Which restaurant would You choose?

Experimentally it was determined that most people choose restaurant which has **more** visitors!

Model of human behavior? (vs Bass diffusion model)




Figure: Incoming new consumers: Bass diffusion model (red curve) against agent based simulation using Kirman herding model (blue curves). Comparison is made for the different number of agents in the system and different discretization periods.

Defining returns



If market is quickly stabilized,

$$D_f + D_c = 0,$$
$$r(t) \approx r_0 \frac{X(t)}{N - X(t)} \Delta \xi(t).$$

One can assume that

the two states in the population dynamics correspond to the chartist trading strategy, excess demand given by

$$D_c = -r_0 X(t) \xi(t),$$

and fundamentalist trading strategy,

$$D_f = [N - X(t)] \ln \frac{P_f}{P(t)}.$$

Stochastic model,

explicitly derived from the previous ABM (one can use birth-death process formalism), for $y = \frac{X}{N-X}$ is given by:

$$dy = \left[\varepsilon_1 + y \frac{2 - \varepsilon_2}{\tau(y)} \right] (1 + y) dt_s + \sqrt{\frac{2y}{\tau(y)}} (1 + y) dW_s,$$

$$\sigma_2 \rightarrow \frac{\sigma_2}{\tau(y)}, \quad h \rightarrow \frac{h}{\tau(y)},$$

$$\varepsilon_i = \frac{\sigma_i}{h}, \quad t_s = ht.$$

The SDE for $y \gg 1$ and assuming that $\tau(y) \sim y^{-\alpha}$ becomes:

$$dy = (2 - \varepsilon_2)y^{2+\alpha}dt_s + \sqrt{2}y^{\frac{3+\alpha}{2}}dW_s.$$

Which has identical form as the general class of SDE reproducing power law statistics (PDF and PSD) [Ruseckas et al., 2011 (Phys. Rev. E)]:

$$dx = \left(\eta - \frac{\lambda}{2} \right) x^{2\eta-1}dt_s + x^\eta dW_s.$$

Comparison yields $\eta = \frac{3+\alpha}{2}$, $\lambda = \varepsilon_2 + \alpha + 1$. Thus we can expect that:

$$p(y) \sim y^{-\varepsilon_2-\alpha-1}, \quad S(f) \sim f^{-\beta}, \quad \beta = 1 + \frac{\varepsilon_2 + \alpha - 2}{1 + \alpha}.$$

Reproducing $1/f$ noise




Figure: Reproducing $1/f$ noise in three cases, $\alpha = 0$ (red squares), $\alpha = 1$ (blue circles) and $\alpha = 2$ (magenta triangles). Other model parameters were set as follows: $\varepsilon_1 = 0.1$, $\varepsilon_2 = 2 - \alpha$. All model data are fitted by: (a) $\lambda = 3$, (b) $\beta = 1$.

The aforementioned general class of SDE

is interesting as it incorporates several widely known stochastic processes into it:

- **Bessel process** ($\eta = 0$): $dx = -\frac{\lambda}{2x}dt + dW$.
- **Squared Bessel process** ($\eta = 0.5$): $dx = \frac{1-\lambda}{2}dt + \sqrt{x}dW$.
- **CIR process** ($\eta = 0.5$, $m = 1$, $x_{min} = 0$, $x_{max} = 1$; the m -exponential diffusion restriction must be applied):
 $dx = \frac{1-\lambda-x}{2}dt + \sqrt{x}dW$.
- **CEV process** ($\eta = \lambda/2$, $m = 2\eta - 2$, $x_{min} = 1$, $x_{max} \rightarrow \infty$; the m -exponential diffusion restriction must be applied):
 $dx = (\eta - 1)xdt + x^\eta dW$.

The same processes should be also reflected by the ABM. Note that CEV and CIR processes can be obtained from the ABM by assuming that $\varepsilon_2 = 2$.




Figure: Numerical results obtained from the CEV-like case, $\alpha = 0$ (red squares), $\alpha = 1$ (blue circles) and $\alpha = 2$ (magenta triangles). Other model parameters were set as follows: $\varepsilon_1 = \varepsilon_2 = 2$.

$$dy = \varepsilon_1 y dt_s + \sqrt{2} y^{\frac{3+\alpha}{2}} dW_s, \quad p(y) \sim y^{-3-\alpha}, \quad S(f) \sim f^{-1-\frac{\alpha}{1+\alpha}}.$$

Variety of reproducible λ and β values





Figure: Wide spectra of obtainable λ and β values. Model parameters were set as follows: $\alpha = 1$, $\varepsilon_1 = 0.1$, $\varepsilon_2 = 0.1$ (red plus), 0.5 (green cross), 1 (blue stars), 1.5 (magenta open squares), 2 (cyan filled squares) and 3 (orange open circles). Black curves correspond to the limiting cases: (a) $\lambda_1 = 2$ and $\lambda_2 = 5$, (b) $\beta_1 = 0.5$, $\beta_2 = 2$

Formulation of the three group agent based model



Transition probabilities for $\vec{X} = \{N_f, N_p\}$,

having in mind that $N_o = N - N_f - N_p$:

$$\begin{aligned}p(\{N_f, N_p\} \rightarrow \{N_f - 1, N_p\}) &= N_f(\epsilon_{cf}/2 + N_o)\Delta t_s, \\p(\{N_f, N_p\} \rightarrow \{N_f + 1, N_p\}) &= N_o(\epsilon_{cf} + N_f)\Delta t_s, \\p(\{N_f, N_p\} \rightarrow \{N_f - 1, N_p + 1\}) &= N_f(\epsilon_{cf}/2 + N_p)\Delta t_s, \\p(\{N_f, N_p\} \rightarrow \{N_f + 1, N_p - 1\}) &= N_p(\epsilon_{cf} + N_f)\Delta t_s, \\p(\{N_f, N_p\} \rightarrow \{N_f, N_p - 1\}) &= N_p(\epsilon_{cc} + HN_o)\Delta t_s, \\p(\{N_f, N_p\} \rightarrow \{N_f, N_p + 1\}) &= N_o(\epsilon_{cc} + HN_p)\Delta t_s.\end{aligned}$$

Stochastic three group model

Stochastic model for $\vec{X} = \{n_f, \xi = \frac{n_o - n_p}{1 - n_f}\}$:

$$dn_f = \frac{1 - 2n_f}{\tau(n_f, \xi)} \varepsilon_{cf} dt_s + \sqrt{\frac{2(1 - n_f)n_f}{\tau(n_f, \xi)}} dW_{s,1},$$

$$d\xi = -\frac{2H\varepsilon_{cc}\xi}{\tau(n_f, \xi)} dt_s + \sqrt{\frac{2H(1 - \xi^2)}{\tau(n_f, \xi)}} dW_{s,2}.$$

Similar equations were obtained in Evolutionary Game Theory from mathematical considerations see [Traulsen et al., 2012 (Phys. Rev. E)].

Fractured spectral density




Figure: Fractured spectral density of various observables. Yet obtained using different parameter sets.

- Nonlinear stochastic model possessing power law spectral density, $S(f) \sim 1/f^\beta$, can be obtained from a microscopic agent based model.
- The nonlinear herding terms in the transition probabilities are essential in reproduction of $1/f$ noise.
- Introducing variability of trading activity generalizes model and offers more modeling possibilities.
- Three group model is able to capture fractured spectral density, but still must be further developed

For further reference see [Kononovicius & Gontis, 2012 (Physica A)], [Ruseckas, Kaulakys & Gontis, 2011 (EPL)].



Physics of Risk
Models of competition and conflicts
in economics



<http://mokslasplus.lt/rizikos-fizika/en>

Thank You!